

STATE OF CALIFORNIA
EARL WARREN, Governor
DEPARTMENT OF NATURAL RESOURCES
WARREN T. HANNUM, Director

DIVISION OF MINES
FERRY BUILDING, SAN FRANCISCO 11
OLAF P. JENKINS, Chief

SAN FRANCISCO

SPECIAL REPORT 5

FEBRUARY 1951

GEOLOGY
OF THE
JURUPA MOUNTAINS
SAN BERNARDINO AND RIVERSIDE COUNTIES,
CALIFORNIA

By EDWARD M. MACKEVETT



GEOLOGY OF THE JURUPA MOUNTAINS SAN BERNARDINO AND RIVERSIDE COUNTIES, CALIFORNIA*

BY EDWARD M. MACKEVETT **

OUTLINE OF REPORT

	Page
Abstract	3
Introduction	3
General geology	4
Pre-batholith metasedimentary rocks	4
Siliceous metamorphic rocks	5
Marble	6
Batholithic rocks	7
San Marcos gabbro	7
Bonsall tonalite	7
Woodson Mountain granodiorite	8
Pegmatites	9
Quaternary surficial deposits	10
Old alluvium	10
Recent deposits	11
Structure	11
Economic geology	12
Marble deposits	12
Granitic rocks	13
Gold and pegmatite prospects	14
References cited	14

Illustrations

	Page
PLATE 1. Geologic map of the Jurupa Mountains	4-5
FIGURE 1. Index map showing location of Jurupa Mountains	4
2. Photomicrograph of quartz-biotite gneiss	5
3. Photomicrograph of quartz-biotite gneiss	5
4. Photomicrograph of quartz-biotite gneiss	6
5. Photomicrograph of quartzite	7
6. Photomicrograph of gabbro	8
7. Photomicrograph of border facies gabbro	8
8. Photomicrograph of gabbro	9
9. Photomicrograph of gabbro	9
10. Photomicrograph of leuco-gabbro	10
11. Photomicrograph of tonalite	10
12. Photomicrograph of granodiorite	11
13. Photomicrograph of granodiorite	12
14. Geologic map of Jensen's quarry	13

ABSTRACT

The Jurupa Mountains are in western San Bernardino and Riverside Counties, California. Their geographical center is about 44 miles east of Los Angeles and 7 miles northwest of Riverside. They are in the northern part of the Perris fault block at the north end of the Peninsular Ranges, which extend southward into Lower California.

Granitic and recrystallized metamorphosed sedimentary rocks make up the greatest part of the Jurupa Mountains. The granitic rocks are the northernmost known exposures of the rocks typical of the southern California batholith, which was probably formed by a series of deep-seated intrusions, not separated by any great period of time but overlapping in age. The intrusions probably came into place by a combination of complex processes of which stoping was the most important.

The oldest rocks in the Jurupa Mountains are a series of metamorphosed sedimentaries which were formed before the granitic intrusions and, in part, were recrystallized and metamorphosed by those intrusions. The resulting rocks consist of quartz-biotite gneiss, quartzite, biotite-quartz schist, marble, various calcareous-siliceous contact rocks, and amphibole schists; this is approximately in the order of their abundance. These remnants of old sedimentary rocks characteristically occur as irregular partitions (septa) between the various granitic intrusions. Jensen's quarry, a locality known to mineral collectors throughout the world for its rare minerals, owes its unusual mineral assemblage to the contact metamorphism brought

about by granitic intrusions into sedimentary rocks. One of the most widely distributed of the darker, more basic of the igneous intrusions is the San Marcos gabbro, which is rich in hornblende and occurs as small island-like bodies within the later intrusive rocks. The Bonsall tonalite, named by Hurlbut, and described and mapped by Larsen in the Peninsular Ranges, is essentially a medium-to coarse-grained granitic rock rather rich in biotite and hornblende. It is marked by abundant dark inclusions near its borders and commonly shows distinct foliation and lineation. It is the most widespread rock in the Jurupa Mountains. The Woodson Mountain granodiorite consists predominantly of light-colored minerals and shows variations in texture. It crops out in large boulder-like masses, and is prominent because of its relatively high resistance to erosion. Pegmatite dikes are abundant in the Jurupa Mountains and form bold, rib-like outcrops. Alluvial fans, locally overlain by wind-blown sand, practically surround the low mountains.

Economic resources of the area are the marble, or metamorphosed limestone, used in making cement, and the several varieties of granitic rocks which are used mainly for rip-rap. In a number of places quartz stringers within the gneiss or schist have been prospected for gold, with relatively little success.

INTRODUCTION

Geography. The Jurupa Mountains are in western San Bernardino and Riverside Counties, California. The center of the mapped area is approximately 44 miles east of Los Angeles and 7 miles northwest of Riverside. Intermittent streams and alluvial fan distributaries trend southward to the Santa Ana River and drain the area. The eastern part of the Jurupa Mountains is characterized by steep ridges that trend about north 15° E.; the western part has moderate slopes and an eastward topographic trend. Maximum relief in the mapped area is 1,490 feet. U. S. Highway 60 traverses the southern part of the mapped region and provides access to the Jurupa Mountains. Vineyards border the mountains on the north, east, and west. Rural residences are scattered throughout the area adjacent to the southern slopes of the Jurupa Mountains.

The climate is semi-arid, and the mountains have a Sonoran type of vegetation characterized by the predominance of chaparral. Nearby Riverside has an average annual rainfall of about 11 inches.

Nature and Scope of Work. This report is based on the mapping of an area of about 15 square miles. The field work was done intermittently from April 1949 to March 1950, and was supplemented from time to time by laboratory investigations. Aerial photographs of the United States Department of Agriculture were used as a base for the mapping, together with enlargements of the United States Geological Survey topographic maps of the Fontana and Guasti quadrangles. Both map and air-photograph bases were on the scale $\frac{1}{4}$ inch equals 1 mile. Most of the work was done on the topographic maps, which proved to be very accurate and detailed.

In addition to the geologic map of the Jurupa Mountains, a plane-table map of Jensen's quarry was made on the scale 1 inch equals 200 feet. Seventy-six rock specimens were collected, and petrographic studies were made of thin sections cut from 15 of these specimens. On the basis of the field and laboratory studies, eight geologic units have been delineated. Limitations of time made

* Prepared in partial fulfillment of the requirements for the degree of Master of Science, Division of Geological Sciences, California Institute of Technology, Pasadena, California. Manuscript submitted for publication July 1950.

** Geologist, U. S. Geological Survey.




FIGURE 1. Index map showing location of Jurupa Mountains.

it impracticable to divide these units further. Moreover, it was not possible to make detailed mineralogical studies of the contact-metamorphic deposits and of the pegmatites in the area.

For purposes of description the Jurupa Mountains are here divided into three parts. (1) a large, nearly isolated hill northeast of Jensen's quarry; (2) the central, more mountainous area from the longitude of Jensen's quarry to the narrows immediately north of Shannahan's quarries; (3) the irregular hills that lie west of Shannahan's quarries.

Purpose. The present investigation was undertaken to obtain a reasonably detailed map of an area previously studied by reconnaissance methods only. This area constitutes one of the northernmost known occurrences of rocks characteristic of the southern California batholith, and it is possible to correlate individual rock types with those so well described from large areas to the south by Larsen.¹ The Jurupa Mountains contain deposits of marble from which cement is made, and also granitic rocks of economic importance. Several of these deposits have been exploited by quarrying methods. Jensen's quarry, a locality renowned for its rare minerals, lies within the mapped area. Pegmatites are abundant, and information on their habits has been obtained by mapping them.

Acknowledgments. The author wishes to express his gratitude to R. H. Jahns of the California Institute of Technology, who suggested and supervised the project, and who provided several valuable suggestions regarding the contents of the manuscript. R. E. Cobb gave his time generously in capably aiding the plane tabling of Jensen's quarry. J. A. Roddick aided in taking the photomicrographs. The Riverside Cement Company was cooperative in permitting geologic investigations at Jensen's quarry.

¹ Larsen, E. S., Batholith and associated rocks of Corona, Elsinore, and San Luis Rey quadrangles, southern California: *Geol. Soc. America Mem.* 29, 1948.

GENERAL GEOLOGY

The Jurupa Mountains are in the northern part of the Perris fault block, as defined by Dudley.² This structural unit is bounded on the east by the San Jacinto fault, on the west by the Elsinore-Chino fault system, and on the north by the base of the San Gabriel Mountains.

The Jurupa Mountains are composed of crystalline rocks and probably contain the northernmost outcrops of igneous rocks that clearly are a part of the southern California batholith.³ These crystalline rocks consist of a complex series of pre-batholith metasedimentary rocks, and batholithic rocks that range from gabbro to leucogranodiorite in composition. Alluvium of the Cucamonga fan, derived from the San Gabriel Mountains to the north, practically embraces the Jurupa Mountains and extends northward for about 10 miles to the base of the San Gabriel Mountains. The San Gabriel Mountains are made up mainly of crystalline rocks and are bounded on the south by a large frontal fault zone. There is a possibility that some of the rocks of the San Gabriel Mountains⁴ are equivalent to some of those in the Jurupa Mountains; however, the writer found little petrographic similarity between the rocks in the two areas, during a cursory examination of some of the San Gabriel rocks.

The nearest exposed rocks west of the Jurupa Mountains are the Tertiary formations of the Puente and San Jose Hills in the vicinity of Pomona. The Santa Ana Mountains, composed of Tertiary and Mesozoic rocks, lie to the southwest. To the east and south, with the exception of local alluvial deposits, the Jurupa Mountains are coextensive with a crystalline rock terrain that is dominated by gabbros, tonalites, and granodiorites of the southern California batholith. The region between the Jurupa Mountains and the base of the San Bernardino Mountains, about 14 miles to the northeast, consists largely of alluvial fans. The San Bernardino Mountains are composed mainly of crystalline rocks.

PRE-BATHOLITH METASEDIMENTARY ROCKS

Scattered bodies of metamorphosed sedimentary rocks that are older than the batholithic intrusions are widespread in the Jurupa Mountains. Gneiss, schist, and quartzite are the most abundantly represented rock types. These have been grouped together on the geologic map (pl. 1) as siliceous metamorphic rocks. Marble is much less common in the Jurupa Mountains, but it has been mapped as a separate unit because of its commercial importance. The degree of metamorphism appears to range from moderate to strong. Local contact zones contain such high-temperature minerals as grossularite, diopside, idocrase, and wollastonite; these indicate high-temperature metamorphic facies near some of the contacts with intrusive rocks. In general, contact metamorphic effects are negligible in the gneiss, schist, and quartzite.


This metamorphic complex has been termed the Jurupa series by Daly,⁵ who believes it to be of questionable Paleozoic age. The Jurupa series probably is cor-

² Dudley, P. H., Physiographic history of a portion of the Perris block, southern California: *Jour. Geology*, vol. 44, p. 358, 1936.

³ Larsen, E. S., op. cit., 1948.

⁴ Miller, W. J., Geology of the western San Gabriel Mountains: *Univ. California at Los Angeles, Pub. in Math. and Phys. Sci.*, 1934.

⁵ Daly, J. W., Paragenesis of the mineral assemblage at Crestmore, Riverside County, California: *Am. Mineralogist*, vol. 20, p. 639, 1935.



GEOLOGIC MAP
OF THE
JURUPA MOUNTAINS

SAN BERNARDINO AND RIVERSIDE COUNTIES
CALIFORNIA

Base enlarged from parts of U.S.
Geological Survey maps of Fontana and Guasti quadrangles

Geology mapped by E.M. MacKevell, 1949-50



Contour interval 25 feet
Datum is mean sea level



Digitized by the Internet Archive
in 2012 with funding from
University of California, Davis Libraries

<http://archive.org/details/geologyofjurupam05mack>




FIGURE 2. Photomicrograph of part of a thin-section of quartz-biotite gneiss. Note parallelism of quartz (q) of similar orientation, and alignment of biotite (b). X nicols. X39

relative with the Triassic (?) Elsinore metamorphic series of Dudley.⁶ The writer prefers to regard metasedimentary rocks of the Jurupa Mountains as being questionably of Triassic age, mainly on the basis of the geographic proximity of the fossiliferous Triassic Bedford Canyon formation⁷ in the Santa Ana Mountains. The Bedford Canyon formation is less metamorphosed, so that the two units are not very similar lithologically. However, the fact that those parts of the Bedford Canyon formation that are of high metamorphic rank strongly resemble the Jurupa Mountain metasediments, and the fact that the Bedford Canyon formation includes lenses of limestone that are similar in size and proportion to the marble lenses in the Jurupa Mountains, suggest a possible correlation.

Woodford⁸ believes that masses of marble at Crestmore, about 2½ miles east of Jensen's quarry, are probably upper Paleozoic in age. He infers this from the occurrence of the fossiliferous Furnace limestone⁹ in parts of the northeastern San Bernardino Mountains, about 40 miles northeast of the Jurupa Mountains, and from evidence cited by Webb¹⁰ that the marble at the Winchester quarry (about 38 miles to the southeast) is Mississippian in age. However, outcrop areas of Furnace limestone are separated from Crestmore (and the Jurupa Mountains) by two major faults of considerable displacement, the San Jacinto and the San Andreas, and Webb's dating is based on only one fossil, a cyathophylloid coral, that was found in a talus pile.

Siliceous Metamorphic Rocks

Siliceous metamorphic rocks constitute the most widespread lithologic unit mapped in the Jurupa Mountains. In the eastern part of the area these rocks are

⁶ Dudley, P. H., Geology of a portion of the Perris block, southern California: California Jour. Mines and Geology, vol. 31, pp. 487-505, 1935.

⁷ Larsen, E. S., op. cit., p. 18, 1948.

⁸ Woodford, A. O., Crestmore minerals: California Div. Mines Rept. 39, p. 336, 1943.

⁹ Woodford, A. O., and Harriss, T. F., Geology of Blackhawk Canyon, San Bernardino Mountains, California: Univ. California, Dept. Geol. Sci. Bull., vol. 17, pp. 265-304, 1928.

¹⁰ Webb, R. W., Evidence of the age of a crystalline limestone in southern California: Jour. Geology, vol. 47, pp. 198-201, 1939.

predominantly biotite-rich gneisses and schists; in the central part biotite-rich gneisses and muscovite- and biotite-bearing quartzites; and in the western part they are mainly impure quartzites. Foliation is commonly well developed in these rocks. In most cases it is steep-dipping and parallels the contact of the nearest intrusive body. In places the foliation is highly contorted and ptygmatically folded. Wherever observed in the same outcrop the foliation and the bedding seem to be parallel in both plan and section. The siliceous metamorphic rocks attain a maximum apparent thickness of about 3,500 feet in the eastern part of the mountains, but the characteristic occurrence of these rocks as narrow septa between masses of intrusive rocks precludes any estimate of original thickness.

Gneiss and Schist. Gneiss and subordinate schist are the most abundant of the siliceous metamorphic rocks. Both the gneiss and the schist contain numerous lit-par-lit stringers, mainly of quartz. The gneissic banding ranges from quartz-rich bands about a millimeter in thickness to bands that attain thicknesses of 2 centimeters. In places the gneiss appears to be intimately associated with material of magmatic derivation.

Most of the gneiss consists chiefly of quartz, subordinate biotite, potash and plagioclase feldspars, and some muscovite. In the typical gneiss gray and dull-white bands alternate on fresh surfaces. The rock as a whole weathers to a light brown.

Less commonly the gneiss is light gray on fresh surfaces and weathers to mottled gray and white. A little of the gneiss is tan on fresh surfaces and brown on weathered ones. Facies containing granulitic textures occur in some of the gneisses. Some of the less-metamorphosed gneisses have local relic textures in part; the subangular shape of many of the minerals in the original sediments is retained. Quartz with a distinct greasy luster is characteristic of a few of the gneisses. Clay minerals, chlorite, and limonite are the principal weathering products of the gneiss.

The schists are less abundant than the gneisses and less diversified in composition. They commonly are gray



FIGURE 3. Photomicrograph of part of thin-section of quartz-biotite gneiss. Field consists mainly of quartz, plagioclase, and biotite. X nicols. X39

biotite-quartz schists that weather dark brown. A dark-gray schist that consists largely of fine-grained amphiboles occurs sparingly throughout the Jurupa Mountains. It is best exposed in the west face of the Decezville quarry where it occurs as inclusions in Bonsall tonalite that reach lengths of several tens of feet.

A typical quartz-biotite gneiss forms a septum between Woodson Mountain granodiorite and Bonsall tonalite on the slopes of the first hill southeast of the eastern Shannahan quarry. It has distinct gneissic banding; the white layers are about a millimeter thick and the gray layers average near 5 millimeters in thickness.

In thin section the gneiss shows an estimated mineral content of one-half to two-thirds quartz, about one-sixth plagioclase, one-tenth biotite, and a remaining tenth composed of augite and orthoclase in nearly equal amounts. Sphene is an abundant accessory, and traces of apatite, opaque grains, and zircon occur as inclusions in biotite. The feldspars, particularly orthoclase, show alteration to clay minerals; the augite and biotite are respectively partly altered to uraltite and chlorite.

A quartz-biotite gneiss collected about a foot from a contact with tonalite near the two adjacent gold prospects in the south-central part of the mapped area consists of felsic bands that reach 15 millimeters in thickness and mafic bands that attain a thickness of 5 millimeters.

Thin-section examination of this phase shows that it is composed of about two-thirds quartz, one-sixth biotite, one-tenth sodic andesine, and one-tenth orthoclase and muscovite. Accessory minerals include sphene, zircon, rutile, and opaque grains. Most of the rock is in grains within the 0.2 of a millimeter to 2.0 millimeter range. The quartz shows undulatory extinction and is slightly fractured. The orthoclase corrodes and commonly encloses plagioclase grains. Alteration products include sericite and clay minerals formed from feldspars, and chlorite formed from biotite.

A thinly banded specimen of quartz-biotite gneiss collected from near the middle of the thick metamorphic section in the eastern part of the area shows the following mineral content when observed beneath a microscope: quartz, one-half to two-thirds; biotite, one-seventh; calcic plagioclase, one-tenth; and muscovite, orthoclase, and hornblende in relatively small fractions. Accessory minerals include sphene, opaque grains, apatite, rutile, and zircon (?). None of the observed quartz grains exceed 0.7 of a millimeter in diameter. These are characteristically sutured and show undulatory extinction. The feldspars are partly altered to clay and sericite. Some of the biotite is altered to chlorite.

Quartzite. Impure quartzite is the prevailing meta-sedimentary rock in the western part of the Jurupa Mountains, where it reaches an estimated maximum apparent thickness of 2,000 feet. The quartzite is a crudely banded, light-gray rock with a greasy, vitreous luster. Weathered surfaces range from yellow brown to dark reddish brown. Biotite and chlorite are the chief constituents of the dark bands in most of the quartzite. Pyrite crystals that reach 0.2 millimeter in diameter are disseminated throughout the quartz of some of the quartzites. Minute needle-like crystals of sillimanite(?) occur locally along fractures within the quartzite. Clay minerals are present in some of the quartzites and probably were




FIGURE 4. Photomicrograph of quartz-biotite gneiss. Field consists mainly of quartz (q) and biotite (b). X nicols. X39

derived from feldspars. Quartzite is resistant to erosion and the trend of many of the ridges in the Jurupa Mountains is controlled by the trend of quartzite beds that underlie them.

A thin-section of typical quartzite, collected in the south-central part of the area, shows that quartz composes from eight-tenths to nine-tenths of the rock. Most of the quartz is in grains that range from 0.2 of a millimeter to 2.0 millimeters in diameter. The remaining fraction contains mostly muscovite, biotite, and hornblende, listed in decreasing order of abundance. Apatite, zircon, opaque grains, and sillimanite were noted as accessory minerals.

Contact-Metamorphic Rocks. Zones of calc-silicate contact rocks that attain 10 feet in thickness are well developed at Jensen's quarry, near the marble bodies a short distance west of Jensen's quarry, and in the westernmost metamorphic rock that appears to be a roof pendant in the Woodson Mountain granodiorite in the western part of the area. These contact rocks are divisible into three general classes: (1) dark-brown, greasy appearing skarn, composed largely of grossularite, generally with subordinate calcite, quartz, and epidote; (2) a white rock that weathers light gray and is composed almost entirely of wollastonite; (3) a quartz-rich rock that ranges from white to light gray in color and is characterized by idocrase, diopside, sphene, grossularite, or augite(?); either singly or in combination.¹¹ Rough surfaces of weathering characterize the outcrops of most of the calc-silicate rocks.

Marble

Recrystallized limestone, chiefly fine- to medium-grained marble, is thickest and most broadly distributed

¹¹ A detailed mineralogical study of the contact-metamorphic deposits is beyond the scope of this report. The mineralogy of the deposits in Jensen's quarry is similar to that at the world-famous localities of nearby Crestmore. Dr. Joseph Murdoch of the University of California at Los Angeles has made some mineralogical investigations at Jensen's quarry and has described some of the rare minerals that occur there. For a detailed description of the mineralogy at Crestmore the reader is referred to Woodford's paper, *Crestmore Minerals* or the paper by J. W. Daly, *Para-genesis of the Mineral Assemblage at Crestmore, Riverside County, California*. Both of these papers are cited in this report and are included in the list of References Cited.

in and around Jensen's quarry, and in the hills immediately west of this quarry. It also occurs at the Glen Avon quarry where it forms definite lenses within siliceous metasedimentary rocks. These lenses are 70 feet in maximum thickness. A bed of marble about a foot thick is present in the westernmost metamorphic body that appears to be a roof pendant in the granodiorite, but it is not shown on the map.

The marble is resistant to erosion and crops out boldly. It is white and is composed largely of medium- to coarse-grained calcite. The marble occurs as beds that rarely exceed 2 feet in thickness. Graphite is abundantly disseminated in a few of these beds, and periclase, brucite, and hydromagnesite also occur in the marble. Rarer constituents, described by Murdoch and Fahey¹² from Jensen's quarry, comprise spinel, pyrite, pyrrhotite, diopside, forsterite, and geikielite. Layers of yellowish predazite, a few inches to a foot thick, are associated with the marble.

The stratigraphic position of the marble in the metasedimentary series is not determinable from evidence in the Jurupa Mountains. Daly¹³ believes that the limestone (marble) deposits at Jensen's quarry are correlative with the Chino limestone (marble) at Crestmore, which he believes to be stratigraphically higher than the greater part of the siliceous metasediments. Commonly the marble is separated from the intrusive rocks by a narrow zone of calc-silicate rock.

BATHOLITHIC ROCKS

San Marcos Gabbro

Isolated bodies of gabbro occur in the southern half of the central and western parts of the Jurupa Mountains. This gabbro is rich in hornblende and is characterized by zoned feldspars in the bytownite range. The gabbro is probably a hornblende-rich facies of the San Marcos gabbro, a widespread member of the southern California

batholith. Miller¹⁴ proposed the name San Marcos gabbro for gabbroic rocks that occur in the San Luis Rey quadrangle. This name has been used loosely to include rocks ranging in composition from norite and olivine gabbro to quartz gabbro. The San Marcos gabbro probably is the oldest rock of the Cretaceous(?) southern California batholith.¹⁵

Two textural varieties of gabbro occur in the Jurupa Mountains. One of these is a dark-gray, medium-grained rock that weathers to a dark reddish brown, and forms the bulk of the gabbro masses. The other is a mottled dark gray and white, brown-weathering rock that is confined to the peripheral zones of the masses. It may represent a facies of the gabbro that was contaminated by later intrusive rocks, although there is no mineralogical evidence to indicate this. The typical texture of the peripheral gabbro consists of zones of medium-grained crystals, surrounded by zones of fine-grained crystals, which is suggestive of a heterogeneous origin for this rock type. Irregular segregations that are rich in hornblende occur in some of the gabbro bodies. Most of the gabbro occupies high topographic positions in the Jurupa Mountains, as it is resistant to erosion; although a few of the gabbro bodies are in areas of subdued topography.

A specimen of gabbro from near the center of the largest gabbro mass in the Jurupa Mountains, a body in the south-central part of the area, is typical of the bulk of the gabbro. It has an average crystal size of about a millimeter in length.

Thin-section studies show it to be composed of about three-fifths plagioclase, one-fifth hornblende, and one-tenth augite. The rock also contains a percent or two each of opaque grains, quartz, and apatite. The plagioclase is zoned in the calcic bytownite range and contains numerous apatite inclusions. The hornblende is partly altered to chlorite, the augite to uraltite, and the plagioclase to clay minerals.

Leucocratic gabbroic facies occur near the margins of some of the gabbro bodies. Thin-section studies show that a specimen of leucogabbro, collected a foot from the eastern contact of the largest gabbro mass in the south-central part of the Jurupa Mountains, is composed of about four-fifths plagioclase, one-tenth hornblende, and less than one-tenth augite. The plagioclase is zoned and ranges in composition from calcic labradorite to sodic bytownite. Most of the plagioclase occurs in grains about 2.0 millimeters long. The hornblende and augite have been partly chloritized, and the augite also has been partly altered to uraltite. Clay minerals have developed from plagioclase.

Bonsall Tonalite

Bonsall tonalite is the most abundant rock in the Jurupa Mountains. This rock was named by Hurlbut¹⁶ from exposures near Bonsall, a village in the San Luis Rey quadrangle in northern San Diego County. This rock is widespread throughout the Peninsular Range areas of southern California and northern Baja California. The Bonsall tonalite is probably correlative with the San Jose quartz diorite of northern Baja California, a rock that

¹² Murdoch, Joseph, and Fahey, J. J., Geikielite, a new find from California: *Am. Mineralogist*, vol. 34, pp. 835-838, 1949.
¹³ Daly, J. W., The geology and mineralogy of the limestone deposits at Crestmore, Riverside County, California: Unpublished Master's Thesis, California Inst. Tech., p. 17, 1931.




FIGURE 5. Photomicrograph of quartzite. Field largely made up of anhedral quartz crystals. X nicols. X39

¹⁴ Miller, F. S., Petrology of the San Marcos gabbro, southern California: *Geol. Soc. America Bull.*, vol. 48, pp. 1397-1426, 1937.

¹⁵ Larsen, E. S., op. cit., p. 45.

¹⁶ Hurlbut, C. S., Jr., Dark inclusions in a tonalite of southern California: *Am. Mineralogist*, vol. 20, pp. 609-630, 1935.



FIGURE 6. Photomicrograph of part of a thin-section of gabbro. Field is dominantly plagioclase (pl) and hornblende (h). X nicols. X39

has been described by Woodford and Harriss.¹⁷ On the basis of field evidence in northern Baja California, Woodford and Harriss believe this rock to be Upper Cretaceous in age.

Typically, the Bonsall tonalite contains numerous inclusions of gabbro, or metamorphic rocks, or both rock types. It varies considerably in texture, and probably represents a series of separate, closely related intrusive masses or plutons. Most of the inclusions are elongate, and few exceed 2 feet in length. Biotite and hornblende, the principal ferromagnesian minerals in the tonalite, commonly are oriented with their long axes parallel. This, together with inclusions, gives much of the rock distinct foliation and lineation. However, some parts of the tonalite lack good foliation and lineation. The Bonsall tonalite is medium to coarse-grained, and ranges in color from light to dark gray.

Salients of tonalite penetrate the gabbro in a few places, and pegmatite dikes genetically related to the tonalite locally cut across the metamorphic rocks. A few of the tonalite-gabbro contacts may be gradational. The contacts between tonalite and metamorphic rocks are in most places nearly vertical. The tonalite disintegrates more readily when exposed to weathering agents than any of the other batholithic rocks. Some deeply weathered areas, largely underlain by tonalite, are well dissected by intermittent streams and gullies and form areas of irregular topography. A good example of this feature occurs on the western slopes of the hills between 1,500 and 2,500 feet southeast of the Declezville quarry. In the north-central part of the Jurupa Mountains, where granodiorite and tonalite are in contact, the granodiorite side of the contact is characterized by large, bouldery outcrops. A subdued, grass-covered surface is on the side underlain by tonalite. Some large residual, boulder-like tonalite bodies protrude above the weathered surface. Weathering "plums," consisting of round bodies of fresh tonalite that reach 4 feet in diameter, apparently entirely surrounded by disintegrated, granular tonalite, are conspicu-

ous in some of the road cuts on Highway 60. Sharp contacts occur between the weathering plums and the disintegrated material. The inclusions are more resistant to weathering, and commonly stand out in bold relief above the tonalite. Thin section examination of tonalite from the mass that underlies the highest peak in the Jurupa Mountains shows the following mineral composition: three-fifths plagioclase feldspar, one-fifth quartz, one-tenth biotite, and a remaining tenth composed principally of hornblende with subordinate augite. Accessory minerals include sphene, zircon, opaque grains, and apatite. The plagioclase is zoned and has an average composition of sodic andesine. The augite is partly altered to antigorite. Sericite and clay minerals have formed at the expense of feldspars, and the biotite and hornblende are partly chloritized.

Woodson Mountain Granodiorite

Granodiorite is best exposed in the vicinity of the second highest peak in the central part of the Jurupa Mountains. This locale consists almost entirely of cliff-like, bold outcrops of Woodson Mountain granodiorite.

The Woodson Mountain granodiorite was named by Miller¹⁸ on the basis of his work in the San Luis Rey quadrangle. The granodiorite is a leucocratic rock that ranges in color from white to light gray and contains only a small proportion of ferromagnesian minerals. It ranges from fine-grained to coarse-grained; the finer-grained facies generally are confined to marginal zones of the plutons. No evidence for the age relationship between the coarse-grained and fine-grained facies was obtained in the Jurupa Mountains. Jahns¹⁹ believes that the fine-grained facies of the Woodson Mountain granodiorite is the younger of the two in the Pala district of southern California, as there it commonly cuts across the coarse-grained granodiorite. The granodiorite is locally gneissic, and in a few places it contains inclusions that appear to have been digested to varying degrees. This

¹⁸ Miller, F. S., op. cit., p. 1399, 1937.

¹⁹ Jahns, R. H., Oral communication, May, 1950.



FIGURE 7. Photomicrograph of part of thin-section of border facies gabbro. Note range in crystal sizes. Plagioclase (pl), hornblende (h), and augite (au) are the dominant minerals. X nicols. X39

¹⁷ Woodford, A. O., and Harriss, T. F., Geological reconnaissance across Sierra San Pedro Martir, Baja California: Geol. Soc. America Bull., vol. 49, pp. 1314, 1331, 1938.




FIGURE 8. Photomicrograph of part of thin-section of gabbro. Note abundant hornblende crystals. X nicols. X39

"mixed zone" is much narrower than that of the Bonsall tonalite and gneissic banding is not as widespread as it is in the tonalite. Rarely the granodiorite has a porphyritic texture, with phenocrysts of flesh-colored microcline, that attain 6 millimeters in maximum dimension, embedded in a fine-grained quartz-feldspar matrix. Contacts between the granodiorite and the metamorphic rocks are nearly vertical.

The Woodson Mountain granodiorite clearly is younger than most of the other crystalline rocks in the area, as it transects them. Many of the pegmatite dikes are derivatives of the Woodson Mountain granodiorite magma, and these cut all the types of crystalline rocks of the Jurupa Mountains. The granodiorite is one of the most resistant of the rocks in the Jurupa Mountains. It commonly crops out in large exfoliation boulders, termed by Larsen ²⁰ "boulders of disintegration." The outer shells of these boulders are moderately disintegrated and crumbly. Small-scale cavernous weathering and incipient pedestal rocks are local features of the granodiorite.

A specimen collected from the westernmost of the twin peaks in the north-central part of the area is a medium- to fine-grained rock. Microscopic examination shows that the specimen is composed of about one-half plagioclase, one-third quartz, one-sixth microcline and less than one-tenth mica, of which biotite is markedly more abundant than muscovite. Accessory minerals include sphene, zircon, and opaque grains. The plagioclase is calcic oligoclase. Most of it occurs in subhedral crystals about 1.0 millimeter long. The quartz grains are anhedral and have a maximum diameter of about 3.0 millimeters. The microcline is subhedral. Both feldspars have partly altered to clay minerals and the plagioclase is also partly sericitized.

Pegmatites

Pegmatite dikes are abundant in the Jurupa Mountains. They are best developed in the eastern and in the western parts of the mapped area. Although pegmatites have not been differentiated on plate 1, three types of pegmatites are represented in the Jurupa Mountains;

(1) pegmatites that appear to be related in composition and genesis to the Bonsall tonalite; (2) pegmatites that appear to be related in composition and genesis to the Woodson Mountain granodiorite; and (3) granite pegmatites that are characterized by a great predominance of potash feldspars and quartz, very large crystals, and irregular areal distribution. The granodiorite pegmatites are the commonest type in the Jurupa Mountains. The granite pegmatites probably are the youngest pegmatites in the area and are the least abundant.

Pegmatites cut across all the other crystalline rock types in the mapped area. The dikes range in width from less than an inch to about 24 feet, and in length from a few feet to at least 5,000 feet. Their bold, rib-like outcrops are conspicuous on the slopes and ridges of the area. The pegmatites have a wide range in crystal size and a variety of subordinate textures. Some of the perthite crystals in the granitic pegmatites exceed 20 centimeters in length. The pegmatites are predominantly white in color, chiefly because of the presence of abundant white perthite.

The tonalite pegmatites are characterized by a large amount of plagioclase. A few of the tonalite pegmatites contain hornblende, for example the westernmost pegmatites shown on plate 1.

Mineralogically, the typical granodiorite pegmatite consists of perthite (generally microcline and oligoclase), light-gray quartz, plagioclase (other than that in the perthite), biotite, black tourmaline, and muscovite. Less common are garnet and limonite (pseudomorphous after pyrite).

The granite pegmatites are characterized by very-coarse crystals, a relatively low plagioclase content, and a more diversified mineralogical assemblage than the other pegmatites.

Murdoch and Webb ²¹ reported beryl crystals from a pegmatite a short distance west of Jensen's quarry. The feldspars are the dominant minerals of most of the pegmatites, but quartz predominates in a few, as in the small

²¹ Murdoch, Joseph, and Webb, R. W., *Minerals of California: California Div. Mines Bull. 136*, p. 70, 1948.



FIGURE 9. Photomicrograph of part of thin-section of gabbro. Large plagioclase crystal has well-developed carlsbad and albite twinning. The plagioclase is almost entirely surrounded by a finer-grained area in which hornblende and augite are the principal minerals. X nicols. X39

²⁰ Larsen, E. S., *op. cit.*, p. 114, 1948.



FIGURE 10. Photomicrograph of part of thin-section of leuco-gabbro, showing abundant large plagioclase crystals. X nicols. X39

bleb-like masses of quartz-rich pegmatite in the central part of the Jurupa Mountains. Some of the pegmatites are locally finer-grained, and aplitic along their outcrops.

Most of the pegmatites have crude symmetrical zoning. Typically, this zoning is as follows: (1) a narrow quartz-rich core commonly with well-developed perthite euhedra and, less commonly, subhedral tourmaline; (2) an intermediate zone, about twice as wide as the core, that consists largely of perthite; this zone may have a graphic texture; biotite is generally abundant contiguous to the core, and muscovite is less commonly present; (3) a wall zone approximately six times the width of the core and consisting mainly of a mixture of medium-grained quartz, feldspar, and biotite. Commonly the wall zone is thinly banded.

There are exceptions to the described internal arrangement for the Jurupa Mountain pegmatites. In places the core lenses out, and the intermediate zone occupies the central position. A few of the pegmatites are asymmetrically zoned, and many are devoid of any clearly recognizable zoning. The phenomenon of multiple zoning, in which several quartz-rich cores are present, occurs in some of the larger pegmatite. Multiple zoning is apparently brought about by the late injection of pegmatitic material into fractures developed in the already solidified portions of the pegmatites.

In general the zoning in the pegmatites of the Jurupa Mountains conforms with the zoning classification proposed by Cameron, Jahns, McNair, and Page;²² the commonest discrepancy being in the characteristics of the wall zone. The wall zones of the Jurupa Mountain pegmatites are generally composed of much smaller crystals than the wall zones typical of the pegmatites of the proposed classification.

Graphic intergrowths are uncommon in the pegmatites of the Jurupa Mountains. They are generally of the well-known graphic granite type, involving perthite and quartz; however, graphic intergrowths between tourmaline and quartz are present in some of the dikes. Some of

the pegmatites that traverse the siliceous metasedimentary rocks contain a few small inclusions of country rock.

The pegmatites resist weathering and are commonly the controlling factor in ridge development. They are most abundant in the siliceous metasediments and may either crosscut or trend parallel to the bedding. Almost all of the pegmatites of the Jurupa Mountains dip steeply. Most of the pegmatites in the western part of the area are not traceable into the granodiorite for more than 2 or 3 feet beyond the granodiorite-siliceous metasediment contact.

The pegmatites of the Jurupa Mountains contrast with most of the famous gem-bearing pegmatites farther south in Riverside and San Diego Counties both in attitude and mineralogy. The Jurupa Mountains pegmatites are much steeper dipping than the typical gem-bearing pegmatites and also lack the lithia minerals that are characteristic of most gem pegmatites.

QUATERNARY SURFICIAL DEPOSITS

Two Quaternary surficial deposits have been differentiated and mapped. They are old alluvium and recent surficial deposits. The recent surficial deposits in the area mapped include: (1) coarse gravel and sand of the Cucamonga fan; (2) aeolian sands; (3) recent alluvium from the Jurupa Mountains; and (4) a few talus and slope-wash deposits which mask significant amounts of the underlying rock.

Old Alluvium

Old alluvium reaches an estimated maximum apparent thickness of 80 feet in the Jurupa Mountains. It is best exposed in the easternmost road cut of Highway 60 within the mapped area. The old alluvium is a crudely stratified fan deposit that consists of sand, gravel, and irregular rock fragments that reach a foot in length. The rocks of the old alluvium were apparently derived from the Jurupa Mountains. This fan material is generally rather well consolidated and occurs at high levels and as bench cappings.



FIGURE 11. Photomicrograph of part of thin-section of tonalite showing relationship between quartz (q), plagioclase (pl), and hornblende (h). X nicols. X39

²² Cameron, E. N., et al., Internal structure of granitic pegmatites: *Econ. Geology*, Mon. 2., p. 20, 1949.

Recent Deposits

Recent surficial deposits are widespread in the mapped area. Their maximum thicknesses were not determined in the present investigation.

Cucamonga Fan Deposits. The Cucamonga fan almost surrounds the Jurupa Mountains. It contains a wide assemblage of clastic material mostly derived from the San Gabriel Mountains. Some of its constituent boulders exceed 2 feet in diameter. Distributaries traversing the Cucamonga fan have locally cut deeply into the fan. Southwest of the Delezville quarry, a long, straight, stream-cut bank, about 40 feet in maximum height, is a conspicuous topographic feature in the alluvium.

Aolian Sands. Aolian sands overlie a large part of the Cucamonga fan in the mapped area. This sand is best exposed along the northern front of the Jurupa Mountains, where the prevailing winds have locally piled it up in drifts and climbing dunes. These dunes and drifts are active, and sand movement can be observed on them on windy days. The constituent sand of these deposits is white in color and consists largely of quartz and feldspars. The aolian deposits probably do not exceed 40 feet in thickness.

Recent Alluvium from the Jurupa Mountains. Recent alluvium from the Jurupa Mountains forms fan deposits bordering most of the southern slopes. This alluvium is characterized by smaller fragments than the typical Cucamonga fan material, and by its content of rocks common in the Jurupa Mountains.

Talus and Slope Wash. Talus and slope wash locally obscure small areas of the underlying rock. Both talus and slope wash are best represented in the vicinity of Jensen's quarry.

STRUCTURE

The Jurupa Mountains may represent a region of structural transition between the prevailing structural trends of the Peninsular Ranges to the south and those of the Transverse Ranges to the north. The general structural trend in the Peninsular Ranges is about N. 35° W. The internal structural trend of the Jurupa Mountains averages N. 70° W.; their topographical trend is east. Both the structural and topographical trends of the San Gabriel Mountains are generally eastward.

The Jurupa Mountains are in the northern part of the Perris fault block. The Perris fault block is bounded by the San Jacinto fault on the east, the frontal fault of the San Gabriel Mountains to the north, and the Losinore-Chino fault system to the west.

Faults. The bulk of evidence in the Jurupa Mountains disfavors any large, continuous faults. The lines of rupture that are mapped as questionable faults on plate are probably essentially joints along which a very small amount of movement took place, and which provided channels for local hydrothermal solutions. These questionable lines of rupture are the only evidence of faulting in the Jurupa Mountains. The presence of faulting is suggested by: (1) a few distinct linear structural features that transect the topography (these are best seen on aerial photographs); (2) a probable local offset and truncation of some of the rock units that strike into the faults; (3) the apparent truncation of a pegmatite dike




FIGURE 12. Photomicrograph of part of thin-section of granodiorite. Large mass of microcline (mi) has numerous poikilitic inclusions. Other minerals in field are quartz (q), biotite (b), and plagioclase (pl). X nicols. X39

by a fault; and (4) local zones of hydrothermally altered rock and siliceous material along the traces of the faults.

Conversely, the only bedrock exposures along the faults reveal no trace of movement, and offset bedding has not been definitely established, mainly because of the dearth of critical outcrops.

It is possible that the Jurupa Mountains are an up-faulted block, and that they are bounded by marginal faults; however, no evidence was found to substantiate this hypothesis and the following negative evidence is offered: (1) the rocks of the margins of the Jurupa Mountains are virtually in the same state of weathering as those that underlie the central parts of the mountains; they are rather fresh and lack shattering, shearing, brecciation, hydrothermal alteration, or other features normally associated with large faults; (2) the borders of the Jurupa Mountains are marked by numerous salients and embayments, and hence do not have the linear aspect that frontal faulting commonly produces.

Internal Structures. Foliation is abundantly developed in the granitic intrusive rocks, especially the tonalite, and in the siliceous metasediments. Generally the foliation in the intrusive rocks and the foliation in the metamorphic rocks are parallel and dip steeply. The foliations are also parallel to the contacts between the two rock types. In most places the contacts dip steeply outward from the intrusive bodies.

Foliation in the intrusive rocks is primary and apparently is due to late-stage magmatic flowage. Evidence for this is the parallel alignment of ferromagnesian minerals and parallelism and elongation of xenoliths. The directional trends of the primary foliation vary; however, the commonest trend is N. 70°-85° W.

Foliation in the metamorphic rocks probably is secondary. This foliation generally trends N. 40°-60° W. in the eastern part of the Jurupa Mountains; its trend is variable in the central part, and in the western part the dominant trend is N. 70°-85° W.

The set of joints best developed in the Jurupa Mountains dips steeply and trends northwest. A second group




FIGURE 13. Photomicrograph of part of thin-section of granodiorite. Crude banding is suggested. Field consists of quartz (q), plagioclase (pl), microcline (mi), and biotite. X nicols. X39

of steeply dipping joints is approximately at right angles to the first, and not as well developed. The Woodson Mountain granodiorite has a set of near-vertical joints approximately normal to its contacts, and a less well-developed set of joints parallel to its contacts. Many other joints in the area, most of them poorly developed, cannot be assigned to any of the above joint sets.

Mode of Emplacement of the Batholithic Rocks. According to Larsen,²³ the primary magma of the batholith was gabbroic in nature, and the different batholithic rock types resulted from the differentiation of this magma. He is of the opinion that stoping probably played an important role in the emplacement of the magma. Evidence of this includes the numerous xenoliths in the different plutons, and parallelism of the contacts between invaded rock and intrusives. In general, evidence in the Jurupa Mountains substantiates Larsen's belief regarding the importance of stoping. Inclusions are plentiful in the intrusive rocks of the area, particularly in the Bonsall tonalite. These inclusions are commonly less than 2 feet in length, but a few attain greater size. The inclusions probably resulted from the incorporation of wall rock in a viscous, cooling magma. The processes of assimilation and forcible intrusion were likely active to a much lesser extent than stoping in the emplacement of the batholithic rocks in the Jurupa Mountains. Assimilation took place on a small scale near many contacts between the intrusive rocks and the siliceous metasediments, and probably on a larger scale at greater depths where hotter magma contacted the invaded rock. Assimilation is suggested by the partially digested condition of some of the inclusions.

Smith²⁴ on the basis of petrofabric analyses of some of the marble from Jensen's quarry states: "The specimens show strong maxima of the optic (C) axes nearly normal to the (intrusive) contact. . . ."

On the basis of girdle patterns at Jensen's quarry, Smith believes that the folding of the marble might have

been caused by the intrusion of the magma. However, this evidence may only be applicable to the latest folding.

Larsen²⁵ believes that the forces that deformed the rocks of the Peninsular Ranges must have acted from the same directions from pre-batholithic times to the present. Daly²⁶ points out that quartz grains in the metamorphic rocks show more evidence of stress than quartz in the igneous rocks, and he believes this to indicate that at least part of the metamorphism had been effected before the intrusion.

It is likely that stoping was the dominant process involved in the emplacement of the Jurupa Mountain intrusive rocks. Assimilation probably took place on a small scale, and forcible intrusion may have occurred locally.

ECONOMIC GEOLOGY

The mineral resources of the Jurupa Mountains that are or have been utilized are: marble, mainly for cement-making purposes, from Jensen's quarry and from the Glen Avon quarry; tonalite from the Declezville quarry; and granodiorite from the Shannahan quarries and from the small unnamed quarry near the 117° 30' meridian. A few smaller-scale operations were active in the past, but none of these were carried beyond the prospecting stage. Several gold prospects are present in the siliceous metasediments and a few in the gabbro. Some of the pegmatites have been prospected by means of small cuts.

Marble Deposits

Jensen's Quarry. Jensen's quarry, where marble is quarried by the Riverside Cement Company for the manufacture of cement, contains the largest and best-known marble deposits in the mapped area. It is in the southeastern part of the Jurupa Mountains. This property is divisible into three main parts: the south quarry, the main quarry, and the north quarry.

The geology at Jensen's quarry is dominated by the intimate association of marble with siliceous metamorphic rocks and salients of intrusive rocks. Bedding in the marble bodies strikes northeast and dips 60°-80° SE. Smith²⁷ believes that the marble at Jensen's quarry may be in a tight, east-plunging fold. However, there does not appear to be any field evidence to substantiate this belief. The marble bodies attain apparent maximum thicknesses of at least 200 feet.

The main quarry is approximately 420 feet long (in a northwesterly direction) and 200 feet wide. The north face of this quarry has a maximum relief of 230 feet. The rim of the south face is not more than 75 feet higher than the quarry floor.

The south quarry is roughly 320 feet in an easterly direction and 200 feet in a northerly direction. The maximum vertical distance between the rims and toes of the faces of this quarry is about 100 feet.

The north quarry is nearly 1,000 feet in length and has a steep southern face that reaches elevations that exceed the elevation of the quarry floor by 150 feet.

The present production is entirely from the main quarry. The irregular association of the marble with other rocks precludes any accurate estimate of the marble

²³ Larsen, E. S., op. cit., p. 171, 1948.

²⁴ Smith, Alexander, Structural petrology at Crestmore, California: Unpublished doctorate thesis (minor), California Inst. of Tech., p. 26, 1947.

²⁵ Larsen, E. S., op. cit., p. 120, 1948.

²⁶ Daly, J. W., op. cit. The geology and mineralogy of the limestone deposits at Crestmore, Riverside County, California, p. 12, 1931.

²⁷ Smith, Alexander, op. cit., p. 16, 1947.

reserves. The hill between the main quarry and the north quarry has been extensively prospected by diamond drilling, and it is likely that the Riverside Company will quarry the marble there at some future time.

Information on the history of Jensen's quarry is rather sketchy. The quarry was apparently developed during World War I by the Riverside Cement Company, and production maintained until 1927, when the property was shut down. There was little activity from 1927 to 1948. The Riverside Company did not resume large-scale operations at Jensen's quarry until 1948. Although no production records are available for Jensen's quarry it is likely supplying a substantial portion of the marble used at Crestmore.

Glen Avon Quarry. The Glen Avon quarry is on the southern slope of the western part of the Jurupa Mountains. The marble occurs in two steeply dipping, lens-like bodies that trend nearly due east. These bodies are within a bedded sequence of siliceous metasedimentary rocks. The marble lenses have maximum exposed dimensions of 750 feet in length and 70 feet in thickness. They have been exploited by two open cuts. The longest of these cuts extends about 150 feet parallel to the strike of the marble bodies. These deposits have not produced for several years. During 1928 two men worked the deposits and shipped marble to Los Angeles, where it was ground and used as chicken grit. The low reserves appear to limit the possibility of future significant production from this deposit.

Other Marble Deposits. A marble body which appears to have commercial possibilities lies about 1,500 feet west of the main Jensen's quarry. Its eastern contact dips 50° E. The only other marble body shown on plate 1 caps a hill about 2,500 feet southwest of the main Jensen's quarry. The apparent small volume of this deposit and its inaccessibility are factors that inhibit future development.

Granitic Rocks

Shannahan Quarries. The Shannahan quarries, where Woodson Mountain granodiorite is obtained, are in the "narrows" near the boundary between the western and central parts of the Jurupa Mountains. The workings consist of two quarries and a few small prospect cuts. The easternmost quarry, the largest of the two, has a face 700 feet in length and 75 feet high. A set of joints, trending almost due east, which are 8 to 10 feet apart, and other joints that trend north, simplify quarrying operations at this location. Weathered debris locally overlies the granodiorite and averages 1 foot in thickness.

These quarries were formerly known as the Bly Brothers quarries. They were first operated around 1900, and granodiorite from them was used for building purposes in Riverside and Los Angeles. In 1928 about 150 tons of granodiorite was produced per month from these workings. The granodiorite was shipped to Los Angeles where it was ground and used for roofing granules and chicken grit. Intermittent operations characterize the recent history of the quarries. Most of the recent production has been used as riprap. In 1949 large quantities of rock obtained at these quarries was used for breakwater construction in the Long Beach harbor. The reserves are large.

Declezville Quarry. The Declezville quarry is in Bonsall tonalite on a northward-projecting salient of the western part of the Jurupa Mountains. It is the oldest and largest quarry in the area. The north-trending face of the Declezville quarry is approximately 1,400 feet in length and 100 feet in maximum height above the quarry floor. The east-trending face is about 700 feet long and rises more than 100 feet above the quarry floor.

The tonalite is relatively free from inclusions. Weathering has formed a dark-brown soil to a depth of about 3 feet below the rim of the quarry. Quarrying is abetted by evenly spaced, northwest-trending joints.

The Declezville quarry is owned by the Southern Pacific Railroad, and was first operated by the railroad prior to 1900. In 1905 practically the entire output was used as rubble in the government breakwater at San Pedro. At the present time the quarry is leased to the Columbia Construction Company, a subsidiary of the Kaiser Steel Corporation. As part of their lease the company produces a small amount of riprap for the Southern Pacific Railroad. The bulk of the recent production was for riprap that was used in the Long Beach breakwater. The quarry has been inactive since the early part of 1950. It is served by a paved road and by a spur line that connects with the main east-west line of the Southern Pacific Railroad. Reserves are vast.

Other Workings in Granitic Rocks. A small working about 50 feet long, 20 feet wide, and 15 feet high is in the Woodson Mountain granodiorite on the northern slope of the Jurupa Mountains near the 117° 30' meridian, at an altitude of 925 feet. The granodiorite at this location is fine-grained and free from inclusions, and probably is quarried for building purposes, although no data are available on the production from this locality. Some other small cuts are in the granodiorite in the western part of the Jurupa Mountains, but none of these are more than prospects.




FIGURE 14. Geologic and topographic map of Jensen's quarry.

Gold and Pegmatite Prospects

Several abandoned gold prospects are present along quartz stringers and blebs within the siliceous metasedimentary rocks. A few prospects are in gabbro, and some of the workings are in tonalite. Aside from a small amount of limonite staining, none of the deposits show any criteria suggestive of ore mineralization.

Most of these deposits appear to be similar to the gold deposits of the Julian schist in the Peninsular Ranges of San Diego County. Both the Jurupa Mountain deposits and the Julian schist deposits generally consist of narrow, irregular quartz stringers in a matrix of siliceous metasedimentary rocks, and the deposits of both of these regions probably were formed by hydrothermal solutions emanating from similar sources.

Two inclined shafts and three prospect adits in a small area on the southern slopes of the central part of the Jurupa Mountains form the greatest concentration of the gold prospects. The longest adit is about 75 feet and the deepest shaft has a slope dimension of approximately 65 feet. The most recent-appearing working in the Jurupa Mountains is in the gabbro on the southern slopes of the western part, and consists of a vertical shaft about 40 feet deep, and has trail and dirt road connections with the highway.

Many of the pegmatites have been prospected by means of small trenches, cuts, and drifts of "coyote hole" proportions. However, there apparently never was any commercial production from these pegmatites.

REFERENCES CITED

- Cameron, E. N., Jahns, R. H., McNair, A. H., and Page, L. R., Internal structure of granitic pegmatites: *Econ. Geology*, Mon. 2, 1949.
- Daly, J. W., The geology and mineralogy of the limestone deposits at Crestmore, Riverside County, California: Unpublished Master's Thesis, California Inst. Tech., 1931.
- Daly, J. W., Paragenesis of the mineral assemblage at Crestmore, Riverside County, California: *Am. Mineralogist*, vol. 20, 1935.
- Dudley, P. H., Geology of a portion of the Perris block, southern California: *California Jour. Mines and Geology*, vol. 31, 1936.
- Dudley, P. H., Physiographic history of a portion of the Perris block, southern California: *Jour. Geology*, vol. 44, 1936.
- Hurlbut, C. S., Jr., Dark inclusions in a tonalite of southern California: *Am. Mineralogist*, vol. 20, 1935.
- Larsen, E. S., Batholith and associated rocks of Corona, Elsinore, and San Luis Rey quadrangles, southern California: *Geol. Soc. America Memoir* 29, 1948.
- Miller, F. S., Petrology of the San Marcos gabbro, southern California: *Geol. Soc. America Bull.*, vol. 48, 1937.
- Miller, W. J., Geology of the western San Gabriel Mountains: *Univ. California at Los Angeles Pub. Math. and Phys. Sci.*, 1934.
- Murdoch, Joseph, and Fahey, J. J., Geikielite, a new find from California: *Am. Mineralogist*, vol. 34, 1949.
- Murdoch, Joseph, and Webb, R. W., Minerals of California: *California Div. Mines Bull.* 136, 1948.
- Smith, Alexander, Structural petrology at Crestmore, California: Unpublished Doctoral Thesis (minor), California Inst. Tech., 1947.
- Webb, R. W., Evidence of the age of a crystalline limestone in southern California: *Jour. Geology*, vol. 47, 1939.
- Woodford, A. O., Crestmore minerals: *California Div. Mines Rept.* 39, pp. 333-365, 1943.
- Woodford, A. O., and Harriss, T. E., Geology of Blackhawk Canyon, San Bernardino Mountains, California: *Univ. California Dept. Geol. Sci. Bull.*, vol. 17, 1928.
- Woodford, A. O., and Harriss, T. E., Geological reconnaissance across Sierra San Pedro Martir, Baja, California: *Geol. Soc. America Bull.*, vol. 49, 1938.

O

